

Outcome of ICAEC 2014

The expected outcomes are as follows;

- To declare importance to prevent pollution by micro-pollutants (declare a statement)
- To make clear environmental pollution by micro-pollutants in South East Asia, East Asia and South Pacific region
- To strengthen relationship of researchers, particularly young researchers between Thailand and Japan

Executive Chair

Takeshi Nakano Osaka University

Advisory Board Members

Kiwao Kadokami	The University of Kitakyushu
Masatoshi Morita	Ehime University
Noriyuki Suzuki	National Institute For Environmental Studies
Susumu Kawahara	Japan Society For Environmental Chemistry
Takumi Takasuga	Shimadzu Techno-Research Inc.
Yoshinori Fujimine	Otsuka Pharmaceutical Co., Ltd

Scientific Committee

Chisato Matsumura	Hyogo Pref. Institute of Environmental Sciences
Daisuke Nakajima	National Institute for Environmental Studies
Daisuke Ueno	Saga University
Haruhiko Nakata	Kumamoto University
Hideshige Takada	Tokyo University of Agriculture And Technology
Hiroshi Yamamoto	The University of Tokushima
Izumi Watanabe	Tokyo University of Agriculture And Technology
Kazuhiko Akutsu	Osaka Pref.Institute of Public Health
Koji Arizono	Prefectural University of Kumamoto
Norihisa Tatarazako	National Institute for Environmental Studies
Seiji Watanabe	National Research And Training Institute, Japan
Shin Takahashi	Ehime University
Shoji Nakayama	National Institute for Environmental Studies
Takanori Sakiyama	Osaka city Institute of Environmental Sciences
Takashi Amagai	University of Shizuoka
Tatsuya Kunisue	Ehime University
Tomohiko Isobe	National Institute for Environmental Studies
Yasuyuki Shibata	National Institute for Environmental Studies
Yoshiharu Shirane	SOWA Consultants Inc.
Yuichi Horii	Center for Environmental Science in Saitama
Amornrat Lertworasirikul	Kasetsart University
Aphichart Rodchanarowan	Kasetsart University
Apirat Laobuthee	Kasetsart University
Chart Chiemchaisri	Kasetsart University
Cheema Sorlalump	Kasetsart University
Daam Settachan	Chulabhorn Research Institute
Gautier Landvot	Kasetsart University

Jutamaad Satayavivad	Chulabhorn Research Institute
Panida Navasumrit	Chulabhorn Research Institute
pongsak noophan	Kasetsart University
Sanya Sirivithayapakorn	Kasetsart University
Wilai Chiemchaisri	Kasetsart University

Japan Committee

Akira Toriba	Kanazawa University
Hatsue Braatz	SheGoTec Japan, Inc.
Hideo Yamazaki	Kinki University
Hiroshi Hoshino	Center for Environmental Science in Saitama
Jun Sekizawa	Communication Center for Food and Health Sciences, Japan
Jun Yoshinaga	The University of Tokyo
Kanji Iwamoto	Nihon BUCHI K.K.
Katsuhiko Nakagawa	Shimadzu Corporation
Kei Nomiyama	Ehime University
Kensaku Kakimoto	Osaka Prefectural Institute of Public Health
Kyoichi Komori	Japan Analytical Instruments Manufacturers' Association
Mari Asami	National Institute of Public Health
Masayuki Kunugi	Kanazawa University
Muneko Nishijo	Kanazawa Medical University
Shinsuke Tanabe	Ehime University
Shunji Hashimoto	National Institute for Environmental Studies
Takeshi Enomoto	JEOL. Ltd.
Takuya Shiozaki	Japan Environmental Sanitation Center
Teruhiko Kido	Kanazawa University
Toru Matsumura	UAE-IDEA Advance Analytical Company Limited
Yasuhiro Hirai	Kyoto University
Yoko Fujikawa	Kyoto University
Yoshitoku Yoshida	Nagoya University
Youko Takagi	Kyoto Electronics Manufacturing Co., Ltd.
Yuji Kunimi	Nagao Clinic

Thai committee

Ministry of Natural Resources and Environmental
 Department of Environmental Quality Promotion
 Environmental Research and Training Center
 Chulabhorn Research Institute
 Pollution Control Department
 Department of Science Service
 Department of Agriculture
 Department of Industrial Works
 Department of Medical Sciences
 Chemical Society of Thailand
 Polymer Society of Thailand
 Kasetsart University
 Center of Excellence on Hazardous Substance Management

C 1-1-1	45	Separation of PCB Isomers #52 and #69 in Kanechlor and Fish Oil Samples Using GC/MS/ECD	Hiroshi TAKAKUWA ¹ , Sadao NAKAMURA ¹ Reiko MATSUMOTO ² , Masahide KAWANO ² , Ichiro TAKEUCHI ² ¹ Agilent Technologies, Japan; ² Faculty of Agriculture, Ehime Univ., Japan
C 1-1-2	38	Selective and Comprehensive Analysis of Organohalogen Compounds by GCxGC-HRTOFMS and MS/MS	Shunji HASHIMOTO ^{1*} , Yasuyuki ZUSHI ¹ , Yoshikatsu TAKAZAWA ¹ , Akihiro FUSHIMI ¹ , Kiyoshi TANABE ¹ , Yasuyuki SHIBATA ¹ ¹ National Institute for Environmental Studies, JAPAN
C 1-1-3	98	Comprehensive GC × GC - MS Analysis of Waste motor Oil from Malaysia reveals <i>in-situ</i> Production of PAHs : A new Pollution Source	Narayanan KANNAN ¹ , Haruhiko MIYAGAWA ² , Riki KITANO ² , Katsuhiko NAKAGAWA ² , Megumi HIROOKA ² , Shunji HASHIMOTO ³ , Vladimir P. BEŠKOSKI ⁴ , Fatemeh A JAHROMI ¹ , Takeshi NAKANO ⁵ ¹ Faculty of Environmental Studies, Universiti Putra Malaysia, Malaysia; ² Shimadzu corporation, Japan; ³ National Institute for Environmental Studies, Japan; ⁴ Faculty of Chemistry, Univ. of Belgrade, Serbia; ⁵ Osaka Univ., Japan
C 1-1-4	87	Advantages of GC/MS/MS Analysis for PCBs, PBDEs and PAHs	Katsuhiko NAKAGAWA ¹ , Kouki TANAKA ¹ , Riki KITANO ¹ , Masato TAKAKURA ¹ , Yuki SAKAMOTO ¹ , Tomoaki KONDO ¹ , Haruhiko MIYAGAWA ¹ ¹ Shimadzu Corporation, Japan
C 1-1-5	55	Screening Analysis of 1200 Micro-Pollutants in Environmental Waters in Vietnam	Kiwao KADOKAMI ^{1*} , Hanh Thi DUONG ^{1,2} , Hong Thi Cam CHAU ¹ , Thao Thanh NGUYEN ² , Trung Quang NGUYEN ² ¹ The Univ. of Kitakyushu, JAPAN; ² Institute of Environmental Technology, Vietnam Academy of Science, Vietnam
C 1-2-1	58	Are PPCPs Really Threats to Ecological Risk?	Hiroshi YAMAMOTO ^{1*} , Norihisa TATARAZAKO ² ¹ The Univ. of Tokushima, Japan; ² National Institute for Environmental Studies, Japan
C 1-2-2	5	Nitro-PAH Product Formation from Heterogeneous Reactions of PAHs with NO ₂ , NO ₃ /N ₂ O ₅ , and OH Radicals : Prediction, Laboratory Studies and Mutagenicity	Narumol JARIYASOPIT ^{1*} , Melissa MC INTOSH ¹ , Kathryn ZIMMERMANN ² , Janet AREY ² , Roger ATKINSON ² , Paul Ha-Yeon CHEONG ¹ , Rich G. CARTER ¹ , Tian-Wei Yu ³ , Roderick H. DASHWOOD ³ , Staci L.M. SIMONICH ^{1,4} ¹ Dept. of Chemistry, Oregon State Univ., USA; ² Air Pollution Research Center, Univ. of California, USA; ³ Institute of Biosciences & Technology, Texas A&M Health Science Center, USA; ⁴ Environmental and Molecular Toxicology, Oregon State Univ., Corvallis, USA.
C 1-2-3	60	Assessment of Phthalate Esters in Aquatic Environment of Pradoo Bay, Rayong Industrial Area, Eastern Thailand	Fairda MALEM ^{1*} , Peerapong SOONTHONDECHA ¹ , Patchara KHAWMODJOD ² , Orapin CHIENHAVORN ² ¹ Environmental Research and Training Center, Thailand; ² Faculty of Science, Kasetsart Univ., Thailand
C 1-3-1	130	Development of the TWO-Vector, Green Fluorescence Protein-Based Bioreporter for BTEX Detection	Naruemon CHUMJAI ^{1,2} , Suwat SOONGLERDSONGPHA ³ , Komkrit SUTTIPONPAMIT ³ , Chatree TANKUNAKORN ³ , Alisa S. VANGNAI ^{2,4*} ¹ International Program in Hazardous Substance and Environmental Management, Graduate School, Chulalongkorn Univ., Thailand; ² Center of Excellence on Hazardous Substance Management, Chulalongkorn Univ., Thailand; ³ Environmental Research and Management Department, PTT Research and Technology Institute, Thailand; ⁴ Faculty of Science, Chulalongkorn Univ., Thailand

COMPREHENSIVE GC × GC - MS ANALYSIS OF WASTE MOTOR OIL FROM MALAYSIA REVEALS *IN-SITU* PRODUCTION OF PAHs: A NEW POLLUTION SOURCE

Narayanan KANNAN¹, Haruhiko MIYAGAWA², Riki KITANO², Katsuhiko NAKAGAWA², Megumi HIROOKA², Shunji HASHIMOTO³, Vladimir P. BEŠKOSKI⁴, Fatemeh A JAHROMI¹, Takeshi NAKANO⁵

¹Faculty of Environmental Studies, Universiti Putra Malaysia, 43400 UPM Serdang, Malaysia; ²Shimadzu corporation, 1, Nishinokyo-Kuwabaracho Nakagyo-ku, Kyoto, Japan; ³National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Japan; ⁴Faculty of Chemistry, University of Belgrade, Studentski trg 1, 11000 Belgrade, Serbia; ⁵Osaka University, 2-4 Yamadaoka, Suita, Osaka, Japan

Key word : PCBs, new source, crankcase waste oil, GC×GC MS, finger printing, environmental forensic

Introduction

Spillage and illegal dumping of waste crankcase oil or leakage of the same from vehicles onto road surfaces with the subsequent washout by street runoff seem to be a continuous source of environmental pollution in an era of vehicular transport. To understand and examine this particular source, a detailed study on the chemical composition of the waste oil is needed. GC-MS is an effective technique in the separation of compounds, as gas chromatography offers good separation and mass spectrometry offers compound specific identification; hence, it is widely used in environmental forensic studies¹. However, crankcase oil is laden with hydrocarbons such as *n*-alkanes; *iso*-alkanes; steranes; terpanes; polynuclear aromatic hydrocarbons (PAHs); aromatic steranes; and specific polycyclic compounds². In this kind of a situation selective detection of target compounds from overlapping co-contaminants is a challenging task. Multidimensional chromatographic techniques have been applied in the determination of polychlorinated biphenyls in complex environmental matrices³⁻⁴. Incorporating this principle in the comprehensive two dimensional GC × GC technique along with mass spectrometry has added ultra-high separation capability to environmental forensic studies⁵⁻⁶. This technology comes handy when separation of target substances from complicated matrices becomes problematic using ordinary GC-MS. We have applied GC × GC - MS technique in the analysis of some waste oil samples from Malaysia and the data were compared with that of GC-MS. Waste motor oil could be an important source of PAHs pollution in Malaysia as vehicular transport is heavy and environmental regulations do not restrict the release of motor oil to the environment strictly. Tropical climate with frequent storms enhance the release of contaminated motor oil to the environment. Application of GC × GC - MS technique has identified and confirmed the *in-situ* production of PAHs through pyrolytic processes in motor engine.

Materials and methods

Waste motor oil (crankcase oil) was sampled at 11 automobile repair shops in the district of Selangor, Malaysia. These were uncharacterized, uncategorized oil samples (MY-1,2,3,4,5,6,9,10,11,12,13) as oil was pooled over a period of time in drums (storage tanks). On the other hand, two crankcase oil samples were collected directly from cars that came for repair (MY-7 & 8). One Castrol unused oil (MY-14) and one Yamaha unused oil (MY-15) were sampled (control) as well. The oil samples were dissolved in iso-octane and analyzed directly in GC-MS and GC × GC - MS.

GC × GC - MS was performed on a gas chromatograph - quadrupole mass spectrometer GCMS-QP2010 Ultra (Shimadzu, Kyoto, Japan) and GC × GC modulator (Zoex Corp.). A DB®-1 (J&W Corp., methyl polysiloxane, 15 m × 0.25 mm I.D., 1.0 µm film thickness) and a Rtx®-WAX (Resteck Corp., 2.5 m × 0.1 mm I.D., df=0.1 µm) columns were connected through the GC × GC modulator as the first and second capillary columns, respectively. Results were processed using a special multipurpose GC × GC analysis software package. It is capable of directly reading GC × GC data obtained with GCMSsolution, converting it to a 2-dimensional image, and then analyzing it.

Results and discussion

The hydrocarbon families in motor lubricating oils are best studied using gas chromatography-mass spectrometry in full scan mode. Various homologous series of hydrocarbons were characterized at known fragment ions. They are: *n*-alkanes; *iso*-alkanes; steranes; terpanes; polynuclear aromatic hydrocarbons (PAHs); aromatic steranes; and specific polycyclic compounds. Used waste lubricating and unused motor lubricating oils

are highly depleted in the paraffinic hydrocarbons (*n*-alkane and *iso*-alkanes)². In spite of that a lot of peaks were detected in a conventional GC-MS analysis which were overlapping in the total ion chromatogram.




Figure 1. a) 2D Chromatogram of MY-04 used oil

b) Thermal modulation of MY-04 used oil

On the other hand with GC × GC - MS system incorporating high-polarity second column, it was possible to separate the paraffin from aromatic series, there by obtaining a blob distribution pattern reflecting the compound's structure. Polyaromatic hydrocarbons (PAHs), such as fluoranthene and pyrene, could be separated from oil matrix and their identification was easily performed using mass spectral similarity search (Fig. 1).

Used oil can be differentiated from unused lubricating oil, because it often has low molecular range hydrocarbons which are absent in the unused oil². The area ratios of phenanthrene to anthracene and the area ratios of fluoranthene to pyrene were useful in evaluating and classifying oil samples. However, while using regular GC-MS technique, anthracene was not detected and the area ratio of fluoranthene/pyrene was lower than




Figure 2. a) Identifying pyrogenic PAHs – GC-MS

b) More clarity in GC×GC-MS (e.g. MY-7&8)

that of the used oils (Fig. 2). The area ratios for thirteen used oil samples obtained from conventional GC-MS analysis did not clarify the differences among the samples. On the other hand, when the same samples were analysed using GC×GC-MS an ultra high resolution separation was obtained and the area ratios calculated based on this analysis differentiated distinctly the used oils, unused oils and oils obtained directly from a car (vs pooled samples of various automobiles over a period of time).




Figure 3. Upper: Phenanthrene identification was marred by alkyl PAHs in a conventional GC-MS system: Lower: GC×GC-MS analysis offered a clear solution to co-elution problems.




Figure 4. Fluoranthene/Pyrene:Phenanthrene/Anthracene ratios of used and unused crankcase oils revealing *in-situ* synthesis of PAHs during combustion. MY-14 is unused oil sample and the rest are used oil samples.

It has been noticed earlier that used crankcase oil carried pyrogenic and petrogenic signatures of PAHs¹. Incorporation of unburned fuel (gasoline or diesel) in the car lead to petrogenic signatures while PAHs generated in the combustion chamber in the engine brought pyrogenic signatures to the crankcase oil. The PAHs profile for “fresh” crankcase oil displayed severe depletion of heavy molecular weight PAHs. Hence the ratio of heavy to light (H/L) molecular weight PAHs had been used to identify pyrogenicity¹. In the same way, fluoranthene and

pyrene were often associated during natural matrices analyses and were considered as typical pyrogenic products derived from high-temperature condensation of lower molecular weight aromatic compounds⁷. In fact, phenanthrene and anthracene are two structural isomers. Because of their different physico-chemical properties, they could behave differently in the environment and could lead to different values for their Phe/An ratio that would give useful information on the PAH origin⁸. Phenanthrene is more thermodynamically stable than anthracene, so, Phe/An ratio is observed to be very high in PAH petrogenic pollution, but lower in pyrolytic contamination cases. Similarly, fluoranthene versus pyrene ratios are used to derive the same conclusion. The current research revealed that conventional GC-MS technique might not reveal the concentrations accurately and hence blur the source identification, whereas, GC×GC-MS analysis will result in unambiguous, accurate and sensitive determination of target compounds even in a complex mixture/matrix. Hence GC×GC-MS technique has identified *in-situ* production of PAHs in automobile combustion processes, thereby, revealing a new source of PAHs pollution to the environment.

Conclusion

In comparison to conventional GC-MS analysis GC×GC-MS analysis revealed higher separation power and accurate determination of target compounds in complex matrix, without any conventional cleanup steps. Thus a complex matrix such as waste motor oil (crankcase oil) when injected directly in GC×GC-MS system was separated into several groups of compounds and allowed accurate determination of target compounds. Such an application in the present study revealed *in-situ* production of PAHs during automobile engine combustion process, thereby revealing a new pollution source of PAHs to the environment. Thus GC×GC-MS could be effectively used in finger printing and source identification in environmental forensic studies.

References:

1. Zakaria MP, Takada H, Tsutsumi S, Ohno K, Yamada J, Kouno E, Kumata H. (2002); *Environ. Sci. Technol.* 36, 1907-1918
2. Kaplan IR, Lu ST, Alimi HM, MacMurphy J. (2001); *Environ. Forensics* 2, 231-248
3. Kannan N, Schulz DE, Petrick G, Duinker JC. (1992); *Intern. J. Environ. Anal. Chem.* 47, 201-215
4. Kannan N, Petrick G, Schulz D, Duinker J, Boon J, van Arnhem, E, Jansen S. (1991); *Chemosphere*, 23, 1055-1076
5. Beens, J, Boelens H, Tijssen R, Blomberg J.(1998); *J. High Resol. Chromatogr.* 21, 47-54
6. Blomberg J, Schoenmakers PJ, Beens J, Tijssen R. (1997); *J. High Resol. Chromatogr.* 20, 539-544
7. Socloa HH, Garriguesb M, Ewald M. (2000); *Mar. Pollut. Bull.* 40(5), 387-396
8. Gschwend PM, Hites RA. (1981); *Geochim. Cosmochim. Acta.* 45, 2359-2367