



"16th European Meeting on Environmental Chemistry"

November 30th – December 3rd 2015

Torino, ITALY

BOOK OF ABSTRACTS

Organized by

Maria Concetta Bruzzoniti and Davide Vione (Department of Chemistry, University of Torino)

Environment Park S.p.A.

On behalf of

ACE - Association of Chemistry and the Environment

Book of Abstracts **"16th European Meeting on Environmental Chemistry"** November 30th – December 3rd 2015 Torino, ITALY

ISBN: 978-88-941168-0-9

Editors: Maria Concetta Bruzzoniti and Davide Vione / Department of Chemistry, University of Torino, ITALY

Printing: Agit Mariogros, Beinasco (TO), ITALY

Design: Luca Rivoira / Department of Chemistry, University of Torino, ITALY

Disclaimer

This book was carefully produced from the author's own submission. Minor formal modifications were made, which in no way interfere with their scientific content. The final printing model was established by the editor according to previous template. Nevertheless, we do not warrant the information contained therein to be free of error. Readers are advised to keep in mind that statement, data, illustrations, procedural details or other items may inadvertently be inaccurate.

No part of this document may be reproduced in any form or by any means without written permissions of the editors.

Scientific Committee

Franco AJMONE MARSAN University of Torino, ITALY Samuel AREY **EPFL,SWITZERLAND** Maria Concetta BRUZZONITI * University of Torino, ITALY **Bogusław BUSZEWSKI** University of Torun, POLAND Josef CASLAVSKY Brno University, CZECH REPUBLIC Luisella CELI University of Torino, ITALY **Gianluigi DE GENNARO** University of Bari, ITALY Massimo DEL BUBBA University of Firenze, ITALY Stuart GIBB University of Highlands and Islands, SCOTLAND Ahmed KETTAB Polytechnic National School, ALGERIA Albert LEBEDEV Moscow State University, RUSSIA **Fabrizio PASSARINI** University of Bologna, ITALY Peter Quinto TRANCHIDA University of Messina, ITALY **Davide VIONE*** University of Torino, ITALY

Organizing Committee

Francesco BARSOTTI Università di Torino, ITALIA Maria Concetta BRUZZONITI University of Torino, ITALY **Davide DAMOSSO** Environment Park S.p.A., Torino, ITALY Fabio GOSETTI University of Piemonte Orientale, ITALY Marco MINELLA University of Torino, ITALY Luca RIVOIRA University of Torino, ITALY **Davide VIONE** University of Torino, ITALY

Conference address

Environment Park Via Livorno 60 – 10144, Torino (IT) www.emec16.com

*chairpersons

P-SOIL - 3

Oil pollution in the vicinity of a heating plant in New Belgrade (Serbia) – influence on the quality of the surrounding soil and sediments

¹Srdjan Miletic, ¹Mila Ilic, ¹Jelena Avdalovic, ¹Tatjana Solevic Knudsen, ¹Vladimir Beskoski, ¹Branimir Jovancicevic, ¹Miroslav Vrvic

¹Institute of Chemistry, Technology and Metallurgy - Department of Chemistry, University of Belgrade, Belgrade (Serbia)

srdjan@chem.bg.ac.rs

Petroleum products have been used for decades in the plants for district heating in Belgrade (Capital of Serbia). The most used derivative for this purpose is a heavy oil fuel. Additionally, raw petrol and ecodiesel are in use as well.

During the months of May and June 2015 an extensive investigation of the pollution of the soil and sediments in the vicinity of a heating plant in New Belgrade (Serbia) was conducted.

According to the geologic and hydro geologic characteristics, the research area is located in the alluvial plains of the Sava River, close to its confluence to the Danube. Because of that, the significance of this research is not only local but also regional.

The soil and sediments were sampled in a zig-zag pattern, from 20 micro locations and 5 different depths making in total 100 of samples.

The results showed that the whole investigated area was contaminated with diesel and a heavy oil fuel which have been used for decades in this heating plant. Although the results did not indicate a significant contamination, the presence of these oil pollutants in the investigated samples indicates that this area should be under continuous monitoring.

OIL POLLUTION IN THE VICINITY OF A HEATING PLANT IN NEW BELGRADE (SERBIA) - INFLUENCE ON THE QUALITY OF THE SURROUNDING SOIL AND SEDIMENTS

Srdjan Miletić¹, Mila Ilić¹, Jelena Avdalović¹, Tatjana Šolević Knudsen^{1,*}, Vladimir P. Beškoski², Branimir Jovančićević² and Miroslav M. Vrvić²

¹Center of Chemistry, Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11001 Belgrade, Serbia ²Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, P.O. Box 158, 11001 Belgrade, Serbia (* corresponding author: tsolevic@chem.bg.ac.rs)

INTRODUCTION

Petroleum products have been used for decades in the plants for district heating in Belgrade (Capital of Serbia). The most used derivative for this purpose is a heavy oil fuel. Additionally, raw petrol and eco-diesel are in use as well.

During the months of May and June 2015 an extensive investigation of the pollution of the soil and sediments in the vicinity of a heating plant in New Belgrade (Serbia) was conducted. This area has a long history of oil pollution due to: NATO bombing of reservoirs (1999), break-down of mazut reservoirs (2009), and accidental spillage from reservoirs over the years.

According to the geologic and hydro geologic characteristics, the research area is located in the alluvial plains of the Sava River, close to its confluence to the Danube (Figure 1). Because of that, the significance of this research is not only local but also regional.



Figure 1, Location of the investigated area



EXPERIMENTAL

The soil and sediments were sampled from 20 micro locations and 5 different depths making in total 100 of samples. Extractable petroleum hydrocarbons were isolated with hexane using a Soxhlet apparatus. The extracts were precleaned on the column packed with Florisil® and analyzed by gas chromatography-mass spectrometry (GC-MS) techniques

An Agilent 7890N gas chromatograph fitted with a HP5-MS capillary column (30 × 0.25 mm, 0.25 μm film; temperature range: 40 °C for 1 min; then 15 °C min⁻¹ to 100 °C for 1 min; then 10 °C min⁻¹ to 310 °C for 15 min and held for 15 min; with helium as the carrier gas (flow rate 1 cm³ min⁻¹) was used. The GC was coupled to a Hewlett- Packard 5972 MSD operated at 70 eV in the 45-550 scan range.

The peaks were identified according to the literature data (Peters et al., 2005., and references therein) or based on the total mass spectra, using mass spectra databases (NIST/EPA/NIH mass spectral library NIST2000, Wiley/NBS registry of mass spectral data, 7th ed., electronic versions.









Figure 5, Gas chromatogram of an extract from a soil sample typical for this locality.

RESULTS

The results showed that the whole investigated area was contaminated with diesel and a heavy oil fuel which have been used for decades in this heating plant. Although the results did not indicate a significant contamination, the presence of these oil pollutants in the investigated samples indicates that this area should be under continuous monitoring.

REFERENCES

Reference Network (Network) (Network

ACKNOWLEDGMENTS

We thank the Ministry of Education, Science and Technological Development of the Republic of Serbia (Projects 176006 and III 43004) for supporting this research.